

UDC 567.3; 551.735(476)

**D. P. Plax**Belarusian National Technical University, 65 Nezavisimosti Ave., 220013 Minsk,  
the Republic of Belarus, agnatha@mail.ru

## ON THE FINDINGS OF THE CARBONIFEROUS ICHTHYOFAUNA IN THE TERRITORY OF BELARUS

In palaeoichthyological terms, the Carboniferous deposits of Belarus have not been specifically studied for a long time. The findings of the ichthyofauna skeletal elements were mostly random and, as a rule, were passed on by geologists to relevant specialists outside Belarus. The results of the study of some of them were published only in a few papers. These few publications provided information on the ichthyofauna stratigraphic distribution in the Carboniferous deposits of Belarus. The author of this article has been engaged in a targeted search and study of the remains of Carboniferous fishes in the country since 2010. Thanks to his targeted research it was possible to significantly replenish the taxonomic composition of the Carboniferous ichthyofauna of Belarus, to clarify its stratigraphic distribution and to provide its detailed description in several papers. It is proposed to include the identified ichthyoassemblages in the regional part of the new stratigraphic chart of the Carboniferous deposits of the territory of Belarus. At present, the research of the remains of the Carboniferous ichthyofauna of Belarus is actively ongoing. Their stratigraphic significance is being clarified for detailed stratification of the Carboniferous deposits of Belarus and for conducting, on their basis, correlations with coeval deposits developed in non-contiguous and far-remote territories.

**Key words:** Belarus; Carboniferous deposits; ichthyofauna; Chondrichthyes; Acanthodii; Actinopterygii; stratigraphy.  
Ref.: 31 titles.

**Д. П. Плакс**Белорусский национальный технический университет, пр-т Независимости, 65, 220013 Минск,  
Республика Беларусь, agnatha@mail.ru

## О НАХОДКАХ КАМЕННОУГОЛЬНОЙ ИХТИОФАУНЫ НА ТЕРРИТОРИИ БЕЛАРУСИ

В палеоихтиологическом отношении каменноугольные отложения Беларуси долгое время специально не изучались. Находки скелетных элементов ихтиофауны носили в основном случайный характер и, как правило, передавались геологами соответствующим специалистам за пределы Беларуси. Результаты изучения некоторых из них были опубликованы только в единичных работах. В этих немногочисленных публикациях были приведены сведения о стратиграфическом распределении ихтиофауны в каменноугольных отложениях Беларуси. Автор данной статьи занимается целенаправленным поиском и изучением остатков каменноугольных рыб на территории страны с 2010 года. Благодаря его целенаправленным исследованиям, удалось существенно пополнить таксономический состав каменноугольной ихтиофауны Беларуси, уточнить ее стратиграфическое распространение и в отдельных работах привести ее детальное описание. Выделенные ихтиокомплексы предлагается внести в региональную часть новой стратиграфической схемы каменноугольных отложений территории Беларуси. К настоящему времени исследование остатков каменноугольной ихтиофауны Беларуси активно продолжается. Выясняется их стратиграфический потенциал для осуществления детальной стратификации каменноугольных отложений Беларуси и проведения на их основе корреляций с одновозрастными отложениями, развитыми на неопредельных и далеко удалённых территориях.

**Ключевые слова:** Беларусь; каменноугольные отложения; ихтиофауна; Chondrichthyes; Acanthodii; Actinopterygii; стратиграфия.

Библиогр.: 31 назв.

**Introduction.** The Carboniferous deposits are distributed within the territory of Belarus in two separate areas: in the southeast — within the Pripyat Trough and in the southwest — within the Volyn Monocline. Within the Pripyat Trough the Carboniferous sediments usually lie with angular

unconformity on the Upper Devonian deposits, and within the Kulazhin Ledge and in the eastern part of the North Pripyat Fault — on the crystalline basement rocks. The Carboniferous deposits are overlapped most often by the Permian or Triassic deposits, less often by the Jurassic, Cretaceous and Paleogene sediments. The stratigraphic completeness and thickness of their section depend on the structural conditions of their occurrence: these deposits are either absent, or represented by the Lower Carboniferous rocks only on raised salt domes and anticlines, the Middle Carboniferous deposits are also developed in synclines and depressions, and the Upper Carboniferous sediments occur there in some places [1].

In the territory of the Volyn Monocline, the Carboniferous deposits lie on rocks of the Lower Devonian — Upper Silurian and are overlain by formations of the Oxfordian Stage of the Upper Jurassic. According to palaeontological data available, the Carboniferous deposits are represented there by sediments of the Tulian, Aleksinian and Mikhailovian Regional Stages of the Viséan Stage of the Lower Carboniferous only [1]. Accordingly, their stratigraphic completeness and thickness are much less than those of the Carboniferous deposits developed within the Pripyat Trough.

The stratigraphy of the Carboniferous deposits of Belarus is mainly justified by the data obtained from the study of the foraminifera, ostracods, brachiopods and spore-pollen complexes [1]. In addition to the above-mentioned orthostratigraphic groups of organisms, some other representatives of fossil organisms are also often found in these deposits: calcareous algae, plant remains, trilobites, bivalves, gastropods, cephalopods, bryozoans, conodonts and fishes. Of the listed organisms, the ichthyofauna only remained unstudied until the mid-1990s. Since 2010 the author engaged in a targeted search and study of the remains of Carboniferous fishes in the country. A comprehensive information on the findings of the Carboniferous ichthyofauna remains currently known in the territory of the country is presented below. A conclusion about their stratigraphic significance for the Carboniferous deposits of Belarus is also suggested.

**Results and discussion.** The earliest reliable information about the findings of the ichthyofauna remains from the Carboniferous deposits was available from the Pripyat Trough. In a number of publications [2—9] devoted to the stratigraphy of the Lower Carboniferous deposits of the Pripyat Trough findings of the ichthyofauna remains were often mentioned together with the other groups of fossil organisms, but without specific Latin names. Only one short message by D. N. Esin, L. A. Petukhova and O. A. Lebedev [10] published in 1995 included for the first time a fish assemblage found in the territory of the Pripyat Trough in the deposits of the Malevkian Regional Stage of the Tournaisian Stage of the Lower Carboniferous, which listed the following taxa: “*Acanthodes*” sp., Ctenacanthidae gen. 2 and Ctenacanthidae gen. 3, *Protacrodus* sp. nov., *Deltopychius armigerus* (Traquair, 1888) [11], *Taeniolepis trautscholdi* (Chabakov, 1927) [12], *Strepheoschema fouldensis* White, 1927 [13], *Aetheretmon valentiacum* White, 1927 [13], *Holurus parki* Traquair, 1881 [14], *Mesopoma pulchellum* (Traquair, 1881) [14], *Styracopterus fulcratus* (Traquair, 1881) [14], *Rhizodus* sp. and Osteolepiformes indet.

An extended collective paper by D. N. Esin and the co-authors [15] dedicated to the Late Devonian and Early Carboniferous vertebrates of the East European Platform was published in 2000. This paper presents the more complete information about the ichthyofauna stratigraphic distribution in the Lower Carboniferous deposits of the Pripyat Trough. The skeletal fish material described in this paper was collected from nine boreholes drilled in the region of the towns of Turov and Petrikov. The palaeoichthyological studies of the rocks from these boreholes allowed the authors to establish the most numerous and taxonomically diverse Malevkian fish assemblage of the Tournaisian Stage in the Pripyat Trough which contained the following taxa: Acanthodii indet., Ctenacanthidae gen. 1, Ctenacanthidae gen. 2, “*Listracanthus*” sp., Ctenacanthidae gen. indet., *Strepheoschema fouldensis* White, 1927 [13], *Taeniolepis trautscholdi* (Chabakov, 1927) [12], *Aetheretmon valentiacum* White, 1927 [13], *Elonichthys* sp. 3, *E.* sp. 4, *Rhadinichthys* sp. 1, Osteolepiformes indet., *Rhizodus* sp. and Rhizodontida indet. According to their data, the fish

assemblage of the Upian Regional Stage of the Tournaisian Stage was less diverse: Ctenacanthidae gen. indet., Osteolepiformes indet., Rhizodontida indet., “*Elonichthys*” cf. *pulcherrimus* Traquair, 1881 [14], *Strepheoschema fouldensis* White, 1927 [13], *Medoevia?* sp., *Holurus parki* Traquair, 1881 [14] and *Rhadinichthys* sp. The vertebrate assemblage established in the rocks of the Gostovian Regional Stage of the Viséan Stage of the Pripyat Trough was quantitatively inferior to the Malevian fish assemblage, but superior to the Upian ichthyoassemblage and was represented by twelve taxa: *Acanthodes* sp., Ctenacanthidae gen. indet., *Protacrodus* sp. nov. 1, “*Listracanthus*” sp., *Aetheretmon valenticum* White, 1927 [13], *Strepheoschema fouldensis* White, 1927 [13], *Holurus parki* Traquair, 1881 [14], *Rhadinichthys* sp., *Elonichthys* sp., Rhizodontida indet., *Styracopterus fulcratus* (Traquair, 1881) [14] and *Mesopoma pulchellum* (Traquair, 1881) [14]. The last two species of fishes were established by D. N. Esin and his co-authors at this stratigraphic level only.

In 2001 O. A. Lebedev in his paper devoted to the palaeontological characteristic of the Middle Carboniferous deposits of the Moscow Syncline [16] mentioned a finding of *Protacrodus sibiricus* Lebedev, 2001 in the Tournaisian-Viséan deposits from the territory of Belarus.

In 2010 the author discovered fish remains represented by a few scales of *Acanthodes* sp., some ctenacanthid type scales and teeth of Actinopterygii gen. indet. [17; 18] in the deposits of the Tulian Regional Stage of the Viséan Stage from the Lelchitsy 16D borehole drilled in the territory of the Pripyat Trough.

In 2010 the author discovered Early Carboniferous fishes in the Komarovka 91z/10 borehole drilled in the Belarusian part of the Volyn Monocline (southwestern part of Belarus) [17—21]. These were found in carbonate-clayey deposits of the Upper Dregovich Subformation of the Dregovich Formation of the Mikhailovian Regional Stage of the Viséan Stage and were represented by rare scales of acanthodians *Acanthodes* sp., single scales and teeth of chondrichthyans Euselachii indet., ctenacanthid and protacrodontid type scales, placoid scale type 1, placoid scale type 2, *Lissodus* sp., Petalodontiformes indet., relatively numerous scales and teeth of actinopterygians *Mesopoma* sp., Actinopterygii gen. et sp. indet. 1, Actinopterygii gen. et sp. indet. 2, Actinopterygii gen. et sp. indet. 3, Actinopterygii gen. et sp. indet. 4, Actinopterygii gen. et sp. indet. 5, Actinopterygii gen. et sp. indet. 6, Actinopterygii gen. et sp. indet. 7, Actinopterygii gen. et sp. indet. 8, Actinopterygii gen. et sp. indet., as well as some indefinable skeletal elements of ray-finned fishes Actinopterygii indet.

In 2018 A. O. Ivanov and D. P. Plax reported some findings of the chondrichthyan scales and teeth in the Devonian and Lower Carboniferous deposits of Belarus [22]. The remains of cartilaginous fishes were taken from the Tournaisian deposits of the Lower Carboniferous of the Pripyat Trough. In particular, the teeth of *Tamiobatis elgae* Ivanov, 2018 [22], *Cladodoides* cf. *wildungensis* (Jaekel, 1921) [23], *Protacrodus* sp., Protacrodontidae indet. and ctenacanthid, hybodontid type scales, *Ohiolepis* type scales, as well as B and C type scales were found in the rocks of the Malevkian Regional Stage. Some teeth of *Tamiobatis* sp., *Cladodoides* cf. *wildungensis* (Jaekel, 1921) [23], *Protacrodus* sp. and ctenacanthid, hybodontid, euselachian type scales, *Ohiolepis* type scales, as well as B and C type scales were found in the deposits of the Upian Regional Stage. A tooth of *Protacrodus* sp. was found in the rocks of the Cherepetian Regional Stage.

Some additional information on cartilaginous fishes from the Famennian deposits of the Upper Devonian, the Tournaisian, Viséan and Serpukhovian sediments of the Lower Carboniferous and the Bashkirian deposits of the Middle Carboniferous was provided in the subsequent publications by A. O. Ivanov and D. P. Plax devoted to the findings of the chondrichthyan remains in the Upper Devonian and Carboniferous deposits of Belarus [24; 25]. In the Tournaisian Stage deposits of the Pripyat Trough, namely, in the rocks of the Malevkian Regional Stage, a finding of a tooth of *Cladodoides* sp., previously unknown findings of teeth of *Thrinacodus ferox* (Turner, 1982) [26], *Danaea* cf. *fournieri* Pruvost, 1922 [27], *Danaea* sp., *Cladodoides* sp., *Protacrodus* sp., *Sphenacanthus* sp., Hybodontiformes indet. A, Hybodontiformes indet. B, ctenacanthid and hybodontid type scales, as well as Chondrichthyes indet. scales in the rocks of the Cherepetian Regional Stage were described in addition to the previously cited chondrichthyan remains [22].

Some teeth of *Reesodus* sp. and Petalodontiformes indet. were found in the deposits of the Dregovich Formation of the Mikhailovian Regional Stage of the Viséan Stage of the Belarusian part of the Volyn Monocline [24; 25]. The following remains were found in the sediments of the Sozhian Regional Stage of the Serpukhovian Stage of the Pripyat Trough: some teeth of *Bransonella lingulata* Ivanov et Ginter, 1996 [28], *Denaëa wangi* Wang, Jin et Wang, 2004 [29], *Cooleyella fordii* (Duffin et Ward, 1983) [30], Helodontiformes indet., buccopharyngeal denticles of Symmoriiformes indet. (*Stemmatias* type) and euselachian type scales [24; 25]. Some ctenacanthid type scales were found in the rocks of the Yastrebian Regional Stage of the Serpukhovian Stage of the Pripyat Trough [24; 25]. And finally, the chondrichthyan remains represented by teeth of *Bransonella nebraskensis* (Johnson, 1984) [31], *Stethacanthus* sp., Squatinactiformes indet., Protacrodontidae indet. B, buccopharyngeal denticles of Symmoriiformes indet. (*Stemmatias* type), ctenacanthid, hybodontid and euselachian type scales were identified in the deposits of the Dvishki Formation of the Pripyatian Regional Stage of the Bashkirian Stage of the Pripyat Trough [24; 25].

Many fish remains represented by the scales of *Acanthodes* sp. and fragments of fin spines of *Acanthodii* gen. indet., scales of *Dipnoi* indet., teeth of *Strepsodus* sp., Onychodontiformes indet., scales of *Rhadinichthys* sp. and Actinopterygii indet. have been recently discovered in the deposits of the Malevkian Regional Stage from several boreholes drilled relatively close to the town of Petrikov (within the Pripyat Trough).

**Conclusion.** To summarize the above, it can be assumed that targeted searches and systematic studies of fishes from the Carboniferous deposits will significantly improve the understanding of their taxonomic composition and permit us to define the more detailed ichthyozones, which can be a reliable tool for the purposes of stratification of these deposits within the territory of Belarus.

The author of the article expresses deep gratitude to V. G. Filippova for assistance in editing the manuscript text in English.

### References

- 1 Golubtsov V. K., Tolstoshev V. I. [Carboniferous system]. *Geology of Belarus*. A. S. Makhnach, et al.; National Academy of Sciences of Belarus, Institute of Geological Sciences; Edited by A. S. Makhnach. Minsk, 2001, 239—259. (in Russian)
2. Kedo G. I. [On the stratigraphy and spore-pollen complexes of the lower horizons of the Carboniferous of the BSSR]. *Reports of the Academy of Sciences of the BSSR*, 1957, vol. 115, no. 6, pp. 1165—1168. (in Russian)
3. Kedo G. I. [The spore-pollen characteristic of the lower horizons of the Carboniferous of the BSSR]. *Transactions of institute for geological sciences*, vol. 1, Minsk, 1958, pp. 46—56. (in Russian)
4. Zingerman A. Ya., Kedo G. I. [On the Devonian and Carboniferous deposits of the southwestern part of the Pripyat Depression]. *Issues of regional geology of the Baltics and Byelorussia*, Riga, 1969, pp. 217—240. (in Russian)
5. Golubtsov V. K. [Carboniferous system]. *Geology of the USSR*, vol. III, Byelorussian SSR. Geological description, Moscow, 1971, pp. 140—161. (in Russian)
6. Kedo G. I., Golubtsov V. K., Demidenko E. K. [Stratigraphy of the Carboniferous deposits of Byelorussia]. *Materials on stratigraphy of Byelorussia*. Minsk, 1981, pp. 69—76. (in Russian)
7. Demidenko E. K. [Biostratigraphic subdivision of the Tournaisian deposits of the western Pripyat Trough based on ostracods]. *Latest data on tectonics, stratigraphy and material composition of sedimentary formations of the BSSR*. Minsk, 1982, pp. 80—88. (in Russian)
8. Avkhimovich V. I., Golubtsov V. K., Demidenko E. K. [Characteristic section of the Devonian-Carboniferous boundary deposits of the Pripyat Depression]. *The Devonian-Carboniferous boundary in the territory of the USSR*. Minsk, 1988, pp. 40—53. (in Russian)
9. Tolstoshev V. I. [Suprasalt Devonian and Carboniferous deposits of the Pripyat Trough]. Minsk, 1988, 150 p. (in Russian)
10. Esin D., Petukhova L., Lebedev O. Vertebrate microremains from the Devonian-Carboniferous deposits of the Pripyat depression (Belarus). *Ichthyolith Issues. Special Publication 1 Socorro*. New Mexico, 1995, p 69.
11. Traquair R. H. Notes on Carboniferous Selachii. *Geological Magazine*, 1888, vol. 5, pp. 81—86.

12. Chabakov A. V. [On crossopterygians from the Russian Carboniferous]. *News of the Geological Committee*, 1927, vol. 46, no. 4, pp. 299—309. (in Russian)
13. White E. I. The Fish-Fauna of the Cementstones of Foulden, Berwickshire. *Trans. Roy. Soc. Edinburgh*, 1927, vol. 55, pp. 255—286.
14. Traquair R. H. Report on fossil fishes collected by the Geological Survey of Scotland in Eskdale and Liddesdale. Part 1, Ganoidei. *Trans. Roy. Soc. Edinburgh*, 1881, vol. 30, pp. 15—71.
15. Esin D., Ginter M., Ivanov A., Lukševičs E., Avkhimovich V., Golubtsov V., Petukhova L. Vertebrate correlation of the Upper Devonian and Carboniferous on the East European Platform. *Courier Forschungsinstitut Senckenberg (Final Report of IGCP 328 project)*, 2000, vol. 223, pp. 341—359.
16. Lebedev O. A. [Vertebrates. *Middle Carboniferous of the Moscow Syncline (southern part)*]. Vol. 2. Palaeontological characteristics. Makhlina, M. Kh. et al. Eds., Moscow, Scientific World, 2001, pp. 92—104. (in Russian)
17. Plax D. P. [Early Carboniferous ichthyofauna of Belarus]. Palaeontology and stratigraphic boundaries. *Proceedings of the LVIII session of the Palaeontological Society of the Russian Academy of Sciences* (April 2—6, 2012, St. Petersburg), St. Petersburg, 2012, pp. 111—112. (in Russian)
18. Plax D. P. Ichthyofauna from the Lower Carboniferous (Viséan) of the Belarusian part of the Volyn Monocline. *Lithosphere*, 2012, no. 1 (36), pp. 3—15.
19. Plax D. P. [On the significance of palaeoichthyological studies for clarifying the boundaries of the modern distribution of the Devonian deposits in the territory of Belarus]. *Vesnik of Brest University. Series 5. Chemistry. Biology. Earth Sciences*, 2016, no. 1, pp. 130—136. (in Belarusian)
20. Plax D. P. [Findings of Palaeozoic ichthyofauna remains in the territory of Belarus]. *BarSU Herald. Series "Biological Sciences (General biology). Agricultural Sciences (Agronomy)"*, 2017, no. 5, pp. 54—64. (in Belarusian)
21. Plax D. P., Kuzmenkova O. F., Obukhovskaya V. Yu., Sachenko T. Ph., Voskoboinikova T. V., Klimentenko Z. M., Bibikova E. V., Karimova L. A. [Lithological and stratigraphic characteristics of Precretaceous deposits of the platform cover of southwestern Belarus (from the results of studying the Komarovka 91z/10 borehole)]. *Lithosphere*, 2012, no. 2 (37), pp. 3—21. (in Russian)
22. Ivanov A. O. & Plax D. P. Chondrichthyans from the Devonian–Early Carboniferous of Belarus. *Estonian Journal of Earth Sciences*, 2018, vol. 67 (1), pp. 43—58.
23. Jaekel O. Die Stellung der Paläontologie zu einigen Problemen der Biologie und Phylogenie. 2. Schädelprobleme. *Paläontologische Zeitschrift*, 1921, vol. 3, pp. 213—239.
24. Ivanov A. O., Plax D. P. Chondrichthyan assemblages from the Upper Devonian — Carboniferous of Belarus. In: "Modern Stratigraphy: methods and applications" and Eighth All-Russian Conference "Upper Palaeozoic of Russia", devoted to the 190th anniversary of the Nikolai A. Golovinsky birth (Kazan, October 28—30, 2024), Kazan, Publishing House of Kazan University, 2024, p. 23.
25. Ivanov A. O. and Plax D. P. New Chondrichthyan Assemblages from the Upper Devonian–Carboniferous of Belarus. *Paleontological Journal*, 2024, vol. 58, Suppl. 4, pp. S391—S402.
26. Turner S. Middle Palaeozoic elasmobranch remains from Australia. *J. Vertebr. Palaeontol.*, 1982, vol. 2, pp. 117—131.
27. Pruvost P. Description de *Denaea fournieri*, sélacien nouveau du Marbre noir de Denée, Part 2. *Bull. Acad. R. Bruxelles*, 1922, ser. 5, no. 8, pp. 213—218.
28. Ivanov A. and Ginter M. Early Carboniferous xenacanthids (Chondrichthyes) from Eastern Europe. *Bull. Soc. Geol. France*, 1996, vol. 167, no. 5, pp. 651—656.
29. Wang N. Z., Jin F. and Wang W. Early Carboniferous fishes (acanthodian, actinopterygians and Chondrichthyes) from the east sector of north Qilian Mountain, China — Carboniferous fish sequence from the east sector of north Qilian Mountain. *Vertebr. Palasiat.*, 2004, vol. 42, pp. 89—110.
30. Duffin C. J. and Ward D. J. Neoselachian shark teeth from the Lower Carboniferous of Britain and the Lower Permian of the U.S.A. *Palaeontology*, 1983, Bd. 26, no. 1, pp. 93—110.
31. Johnson G. D. A new species of *Xenacanthoides* (Chondrichthyes, Elasmobranchii) from the Late Pennsylvanian of Nebraska. In *Papers in Vertebrate Paleontology Honoring Robert Warren Wilson*. Mengel R. M. Ed. *Carnegie Mus. Nat. Hist., Spec. Publ.*, 1984, no. 9, pp. 178—186.

#### Список цитируемых источников

1. Голубов, В. К. Каменноугольная система / В. К. Голубов, В. И. Толстошеев // Геология Беларуси / Ин-т геол. наук ; под. общ. ред. А. С. Махнач [и др.]. — Мн., 2001. — С. 239—259.
2. Кедо, Г. И. О стратиграфии и спорово-пыльцевых комплексах нижних горизонтов карбона БССР / Г. И. Кедо // Доклады Академии наук БССР. — 1957. — Т. 115, № 6. — С. 1165—1168.
3. Кедо, Г. И. Спорово-пыльцевая характеристика нижних горизонтов карбона БССР / Г. И. Кедо // Труды института геологических наук. — Мн., 1958. — Вып. 1. — С. 46—56.

4. Зингерман, А. Я. О девонских и каменноугольных отложениях юго-западной части Припятской впадины / А. Я. Зингерман, Г. И. Кедо // Вопросы региональной геологии Прибалтики и Белоруссии. — Рига, 1969. — С. 217—240.
5. Голубцов, В. К. Каменноугольная система / В. К. Голубов // Геология СССР. — М., 1971. — Т. III : Белорусская ССР. Геологическое описание. — С. 140—161.
6. Кедо, Г. И. Стратиграфия каменноугольных отложений Белоруссии / Г. И. Кедо, В. К. Голубов, Э. К. Демиденко // Материалы по стратиграфии Белоруссии. — Мн., 1981. — С. 69—76.
7. Демиденко, Э. К. Биостратиграфическое расчленение турнейских отложений запада Припятского прогиба по остракодам / Э. К. Демиденко // Новейшие данные по тектонике, стратиграфии и вещественному составу осадочных образований БССР. — Мн., 1982. — С. 80—88.
8. Авхимович, В. И. Характерный разрез пограничных отложений девона и карбона Припятской впадины / В. И. Авхимович, В. К. Голубцов, Э. К. Демиденко // Граница девона и карбона на территории СССР. — Мн., 1988. — С. 40—53.
9. Толстошеев, В. И. Надсолевые девонские и каменноугольные отложения Припятского прогиба / В. И. Толстошеев. — Мн., 1988. — 150 с.
10. Esin, D. Vertebrate microremains from the Devonian-Carboniferous deposits of the Pripyat depression (Belarus) / D. Esin, L. Petukhova, O. Lebedev // Ichthyolith Issues. Special Publication 1 Socorro. — New Mexico, 1995. — P. 69.
11. Traquair, R. H. Notes on Carboniferous Selachii / R. H. Traquair // Geological Magazine. — 1988. — Vol. 5. — P. 81—86.
12. Хабаков, А. В. О кистеперых из русского карбона / А. В. Хабаков // Известия Геологического комитета. — 1927. — Т. 46, № 4. — С. 299—309.
13. White, E. I. The Fish-Fauna of the Cementstones of Foulden, Berwickshire / E. I. White // Trans. Roy. Soc. Edinburgh. — 1927. — Vol. 55. — P. 255—286.
14. Traquair, R. H. Report on fossil fishes collected by the Geological Survey of Scotland in Eskdale and Liddesdale. Part 1, Ganoidei / R. H. Traquair // Trans. Roy. Soc. Edinburgh. — 1881. — Vol. 30. — P. 15—71.
15. Vertebrate correlation of the Upper Devonian and Carboniferous on the East European Platform / D. Esin, M. Ginter, A. Ivanov, E. Lukševičs // Courier Forschungsinstitut Senckenberg (Final Report of IGCP 328 project). — 2000. — Vol. 223. — P. 341—359.
16. Лебедев, О. А. Позвоночные / О. А. Лебедев // Средний карбон Московской синеклизы (южная часть). — М., Науч. мир, 2001. — Т. 2 : Палеонтологическая характеристика. — С. 92—104.
17. Плакс, Д. П. О раннекаменноугольной ихтиофауне Беларуси / Д. П. Плакс // Палеонтология и стратиграфические границы : материалы LVIII Сессии палеонтол. о-ва при РАН (2—6 апр. 2012 г., Санкт-Петербург). — СПб., 2012. — С. 111—112.
18. Plax, D. P. Ichthyofauna from the Lower Carboniferous (Viséan) of the Belarusian part of the Volyn Monocline / D. P. Plax. — Lithosphere. — 2012. — No. 1 (36). — P. 3—15.
19. Плакс, Дз. П. Аб значэнні палеаіхтыялагічных даследаванняў для ўдакладнення межаў сучаснага распаўсюджвання дэвонскіх адкладаў на тэрыторыі Беларусі / Дз. П. Плакс // Веснік Брэсцкага ўніверсітэта. Серыя 5: Хімія. Біялогія. Навукі аб Зямлі. — 2016. — № 1. — С. 130—136.
20. Плакс, Дз. П. Аб знаходках рэшткаў палеазойскай іхтыяфаўны на тэрыторыі Беларусі / Дз. П. Плакс // Вестник БарГУ. Серия «Биологические науки (общая биология). Сельскохозяйственные науки (агрономия)». — 2017. — № 5. — С. 54—64.
21. Литолого-стратиграфическая характеристика домеловых отложений платформенного чехла юго-западной Беларуси (по результатам изучения скважины Комаровка 913/10) / Д. П. Плакс, О. Ф. Кузьменкова, В. Ю. Обуховская [и др.] // Літасфера. — 2012. — № 2 (37). — С. 3—21.
22. Ivanov, A. O. Chondrichthyans from the Devonian–Early Carboniferous of Belarus / A. O. Ivanov, D. P. Plax // Estonian Journal of Earth Sciences. — 2018. — Vol. 67 (1). — P. 43—58.
23. Jaekel, O. Die Stellung der Paläontologie zu einigen Problemen der Biologie und Phylogenie. 2. Schädelprobleme / O. Jaekel // Paläontologische Zeitschrift. — 1921. — Vol. 3. — P. 213—239.
24. Ivanov, A. O. Chondrichthyan assemblages from the Upper Devonian — Carboniferous of Belarus / A. O. Ivanov, D. P. Plax // Восьмая Всерос. конф. «Верхний палеозой России». Современ. стратиграфия: методы и приложения, посвящ. 190-летию со дня рождения Н. А. Головкинского (28—30 окт. 2024 г., Казань (Россия)). — Казань, 2024. — С. 23.
25. Ivanov, A. O. New Chondrichthyan Assemblages from the Upper Devonian–Carboniferous of Belarus / A. O. Ivanov, D. P. Plax // Paleontological Journal. — 2024. — Vol. 58, Suppl. 4. — P. S391—S402.
26. Turner, S. Middle Palaeozoic elasmobranch remains from Australia / S. Turner // J. Vertebr. Palaeontol. — 1982. — Vol. 2. — P. 117—131.
27. Pruvost, P. Description de Denaea fourrieri, sélacien nouveau du Marbre noir de Denée, Part 2 / P. Pruvost // Bull. Acad. R. Bruxelles — 1922. — Ser. 5, no. 8. — P. 213—218.
28. Ivanov, A. Early Carboniferous xenacanthids (Chondrichthyes) from Eastern Europe / A. Ivanov, M. Ginter // Bull. Soc. Geol. France — 1996. — Vol. 167, no. 5. — P. 651—656.

29. Wang, N. Z. Early Carboniferous fishes (acanthodian, actinopterygians and Chondrichthyes) from the east sector of north Qilian Mountain, China — Carboniferous fish sequence from the east sector of north Qilian Mountain / N. Z. Wang, F. Jin, W. Wang // *Vertebr. Palasiat.* — 2004. — Vol. 42. — P. 89—110.

30. Duffin, C. J. Neoselachian shark teeth from the Lower Carboniferous of Britain and the Lower Permian of the U.S.A. / C. J. Duffin, D. J. Ward // *Palaeontology.* — 1983. — Bd. 26, no. 1. — P. 93—110.

31. Johnson, G. D. A new species of *Xenacanthoides* (Chondrichthyes, Elasmobranchii) from the Late Pennsylvanian of Nebraska / G. D. Johnson // In *Papers in Vertebrate Paleontology Honoring Robert Warren Wilson*. Mengel R. M. Ed. Carnegie Mus. Nat. Hist. — Spec. Publ. — 1984. — No. 9. — P. 178—186.

Received by the editorial staff 22.02.2026.